

The porous atmosphere of η -Carinae

Nir J. Shaviv

*Canadian Institute for Theoretical Astrophysics, University of Toronto
60 St. George St., Toronto, ON M5S 3H8, Canada*

ABSTRACT

We analyze the wind generated by the great 20 year long super-Eddington outburst of η -Carinae. We show that using classical stellar atmospheres and winds theory, it is impossible to construct a consistent wind model in which a sufficiently *small* amount of mass, like the one observed, is shed. One expects the super-Eddington luminosity to drive a thick wind with a mass loss rate substantially higher than the observed one. The easiest way to resolve the inconsistency is if we alleviate the implicit notion that atmospheres are homogeneous. An inhomogeneous atmosphere, or “porous”, allows more radiation to escape while exerting a smaller average force. Consequently, such an atmosphere yields a considerably lower mass loss rate for the same total luminosity. Moreover, all the applications of the Eddington Luminosity as a strict luminosity limit should be revised, or at least reanalyzed carefully.

To appear in the Astrophysical Journal Letters

Subject headings: Radiative transfer — hydrodynamics — instabilities — stars: atmospheres — stars: individual (η Carinae)

1. Introduction

η -Carinae is probably one of the most remarkable stellar object to have ever been documented. About 150 years ago, the star began a 20 year long giant eruption during which it radiated a supernova-like energy of roughly 3×10^{49} *ergs* (Davidson & Humphreys 1997). Throughout the eruption it also shed some $1 - 2 \mathcal{M}_{\odot}$ of material carrying approximately 6×10^{48} *ergs* as kinetic energy (Davidson & Humphreys 1997), while expanding at a velocity of 650 *km/sec* (Hiller & Allen 1992, Currie et al. 1996). η -Carinae can therefore serve as a good laboratory for the study of atmospheres at extreme luminosity conditions.

At first glance, it appears that the star shed a large amount of material. Indeed, the inferred mass loss rate during the great eruption of $\sim 0.1 \mathcal{M}_{\odot}/yr$ is significantly larger than the mass loss rate inferred for the star today ($\lesssim 10^{-3} \mathcal{M}_{\odot}/yr$, Davidson & Humphreys 1997 and references therein). However, considering that the luminosity during the great eruption is estimated to be significantly above the Eddington limit, we shall show that the star should have had a much higher

mass loss rate. In fact, it should have lost during the 20 year eruption more mass than its total mass, giving rise to an obvious discrepancy.

A review of our current knowledge of η Car can be found in Davidson & Humphreys (1997). In section 2 we summarize how a wind solution for the star η Car should be constructed. Since the luminosity is very high, the effects of convection must be taken into account. In section 3 we integrate the wind equations to show that no consistent solution for η Car exists within the possible range of observed parameters. Section 4 is devoted to possible classical solutions to the discrepancy, showing that no such possibility exists. In section 5, we show that a porous atmosphere is a simple and viable solution to the wind discrepancy.

2. Solving for the Wind

Since the mass of η -Car is estimated to be of order $100\text{--}120\,M_{\odot}$ (Davidson & Humphreys 1997), the average luminosity in the great eruption was clearly super-Eddington (of the order of 5 times the Eddington limit). That is to say, the radiative force upwards, assuming the smallest possible opacity (for ionized matter) given by Thomson scattering, was significantly larger than the gravitational pull downwards. Optically thin winds formally diverge at the Eddington limit (e.g., Kudritzki et al. 1989 and references therein). Consequently, a consistent wind solution requires an optically thick wind. We thus look for a wind in which the sonic point (which is the point at which the local speed of the outflow equals the speed of sound) is below the photosphere. Moreover, since the duration of the eruption is longer than the sound crossing time of the star by about a factor of 50, a stationary wind appears to be a good approximation.

In practically all super-sonic wind theories which describe super-sonic outflows from an object at rest, a consistent stationary solution is obtained only when the net driving force of the wind (excluding the pressure gradient) vanishes at the sonic point¹. Thus, material experiencing a super-Eddington flux necessarily has to be above the sonic point. If most of the envelope carries a super Eddington flux, then no consistent stationary wind solution can be obtained and in fact, the object will evaporate on a dynamical time scale. In most systems however, this need not be the case. For example, in very hot systems (e.g., hot neutron stars during strong X-ray bursts, Quinn & Paczynski 1985), the opacity in the deep layers is reduced due the reduced Klein-Nishina opacity for Compton scattering at high temperatures. Thus, the sonic point in these objects is found where the temperature is high enough to reduce the opacity to the point where the flux corresponds to the local Eddington limit.

¹The exception is line driven winds in which the force is explicitly a function of dv/dr which is actually an approximation to the line transfer equations. If we had written the proper radiation transfer equations for this case which only *implicitly* depend on dv/dr , we would have recovered that the sonic point coincides with the point at which the total force vanishes (cf Mihalas & Weibel Mihalas 1984 §107). Moreover, line driven winds are important only under optically thin conditions while we describe the optically thick part of the wind.

Another important effect, which should be taken into account, is convection. Deep inside the atmosphere, convection can carry a significant part (or almost all) of the energy flux, thus reducing the radiative pressure to a sub-Eddington value. In fact, as the radiative flux approaches the Eddington limit, convection generally arises and carries the lion share of the total energy flux (if it can) to keep the system at a sub-Eddington level (Joss et al. 1973). Although the total flux in the entire envelope (or almost all of it) can be equivalent to a super-Eddington flux, up to some depth below the photosphere, convection carries most of the flux so as to reduce the radiative flux alone into a sub-Eddington value. A consistent wind solution should therefore, have its sonic point at the location where the most efficient convection cannot carry enough flux any more. As we shall soon see, the problem in η -Car is that this point is relatively deep within the atmosphere, where the density is so high that the expected mass loss is significantly *higher* than the observed one.

To see this in a robust way we integrated numerically the wind equations starting from the photosphere inwards. The equations are those that describe optically thick spherically symmetric winds (Quinn & Paczynski 1985; Żytkow 1972; Kato & Hachisu 1994). The equations of mass conservation, momentum conservation and temperature gradient are

$$4\pi r^2 \rho v = \dot{M} = \text{const} \quad (1)$$

$$v \frac{dv}{dr} + \frac{GM}{r^2} + \frac{1}{\rho} \frac{dP_g}{dr} - \frac{\chi \mathcal{L}_r}{4\pi r^2 c} = 0 \quad (2)$$

$$\frac{dT}{dr} = -\frac{3\chi\rho\mathcal{L}_r}{16\pi r^2 c a T^3} \left(1 + \frac{2}{3\chi\rho r}\right), \quad (3)$$

with standard notation. The parenthesized term in the last equation is a simple approximate interpolation that has the correct asymptotic limits for optical depths much larger and much smaller than unity (Quinn & Paczynski 1985). The last equation is the integrated form of the energy conservation equation. Unlike the aforementioned references, we specifically include advection by a maximally efficient convection. Thus, the integrated form of the energy conservation equation becomes

$$\mathcal{L}_r + \dot{M} \left(\frac{v^2}{2} + w - \frac{GM}{r} \right) + \mathcal{L}_{conv} \equiv \quad (4)$$

$$\mathcal{L}_r - \frac{GM\dot{M}}{r} + \mathcal{L}_{adv} + \mathcal{L}_{conv} = \Lambda_{tot} = \mathcal{L}_{obs} + \mathcal{L}_{kin,\infty}, \quad (5)$$

where Λ_{tot} , $\dot{M}$, $\mathcal{L}_{obs}$ and $\mathcal{L}_{kin,\infty}$ are the total energy output of the star, the wind mass loss rate, the observed luminosity at infinity and the kinetic energy flux at infinity. On the other hand, $\mathcal{L}_r, v, w, \mathcal{L}_{conv}, \mathcal{L}_{adv}$ are respectively, the local radiative luminosity, velocity, enthalpy, convective flux and advected flux (as internal and kinetic energies). The expression adopted for $\mathcal{L}_{conv}$ is $4\pi r^2 u v_s$ where u is the internal energy per unit volume and v_s is the speed of sound. By no means can convection be more efficient than this expression since highly dissipative shocks are unavoidable at higher speeds. It is likely that the maximally efficient convection is somewhat less efficient than this expression, but this will only aggravate the problem that we shall soon expose. Detailed

calculations of the wind were carried out. The calculations include the latest version of the OPAL opacities (Iglesias & Rogers 1996). It is found that the total opacity below the photosphere has comparable contributions from Thomson scattering and absorption processes. This implies that the *modified* Eddington limit, in which the Thomson opacity is replaced by the local total opacity, is somewhat lower than the classical Eddington limit.

Since $T_{eff} \sim 9000^\circ\text{K}$ ², the average luminosity implies a photospheric radius of 10^{14} cm. Note that since it is a thick wind, the exact definition of the photosphere is ambiguous. Nevertheless, different definitions do not change the results by more than 10 – 20%. Knowing that the observed mass loss rate is roughly $0.1 \mathcal{M}_\odot/\text{yr}$ (which gives the observed $2 \mathcal{M}_\odot$ of shed material in 20 years, Davidson & Humphreys 1997), a specified flow speed at the photosphere can be translated to a required density. We can therefore integrate the wind equations inwards. If a consistent wind solution can be obtained for some value of the imposed velocity in the photosphere (which has to be between v_s and v_∞), then the integration inwards should reach a sonic point at which the total force on the gas vanishes. This will be attained if the convective and advective fluxes can carry a significant amount of the total flux so as to reduce the residual radiative flux to a sub-Eddington one.

3. The Discrepancy

We define the luminosity needed to be carried by convection and advection in order to bring about the vanishing of the total local force as $\mathcal{L}_{crit}$. If enough energy is advected and convected then $\mathcal{L}_r$ will be reduced to the local modified Eddington flux:

$$\mathcal{L}_{Edd,mod} = \frac{4\pi cGM}{\chi} \quad (6)$$

with χ the local opacity which can be larger than the Thomson opacity. Thus, from eq. (5), the critical advective+convective flux can be written as

$$\mathcal{L}_{crit} = \Lambda_{tot} + \frac{GM\dot{M}}{r} - \mathcal{L}_{Edd,mod}. \quad (7)$$

Figure 1 shows the fraction $\eta \equiv (\mathcal{L}_{adv} + \mathcal{L}_{conv})/\mathcal{L}_{crit}$ at the sonic point. A consistent solution can be found only if $\eta = 1$ at the sonic point.

Inspection of the figure clearly shows that the space of possible observed values does not contain a viable and consistent solution. This is of course irrespective of whether a solution from the

²This is the typical observed effective temperature for LBV's during outbursts (Humphreys & Davidson 1994). If the temperature is higher than this value, the inferred bolometric magnitude of η Car during the eruption would be more negative, increasing the Eddington factor. If the temperature is lower than $\sim 7000^\circ\text{K}$, the opacity at the photosphere and outwards rises abruptly (Davidson 1987), thus reducing the modified Eddington limit. In both cases, the discrepancy will be aggravated.

photosphere outward can or cannot be obtained. The discrepancy arises because a wind corresponding to the observed low mass loss rate necessarily has a sonic point that is not deep enough to have either convection or advection as an efficient mean of transporting energy. This can be seen from the optical depth at which the sonic point is obtained. In all cases, $1 \lesssim \tau < 300$. However, convection is efficient only up to an optical depth of $\tau \sim c/v_s \gg 300$ for $p_{rad} \sim p_{gas}$ (Shaviv 2000b), or even deeper for larger radiation pressures (i.e., when close to the Eddington limit).

4. Unfeasible Solutions to the Discrepancy

Can the discrepancy be resolved with a classical assumption? Since the discrepancy is rather large, assuming the wind to emerge from an angular fraction f from the star does not relax the problem (it actually aggravates the problem because more material will be blown away from the higher luminosity regions). Another possibility that fails is having a higher velocity in the photosphere than the one observed today for the shed material. This might be the case if the wind collides with previously ejected slow moving material. Even if such material did exist, the necessarily reduced mass loss rate inferred from the present day observed momentum aggravates the problem.

The problem is not mitigated if we relax the assumption that the mass loss rate and the luminosity are assumed to be constant in time throughout the eruption.

If one wishes to solve the problem using magnetic fields, then a solution can be found only if the magnetic energy density at the photosphere is significantly larger (by several orders of magnitude) than the equipartition value with the gas pressure. This of course seems unlikely.

Another option is to have the distance estimate to η -Car be three times smaller than 2300 pc. A shorter distance will remove η -Car out of the cluster Tr16 of massive stars inside which it is observed and leave it instead roaming the inter galactic arm space. Considering the short lifetime of the star, just a few million years, this possibility appears as very unlikely.

The problem can be solved if the mass of the star corresponds to a sub-Eddington luminosity. This proposed solution requires η -Car to be at least a $\sim 1000 M_{\odot}$ star. However, this suggestion is at variance with much lower estimates (see for instance Davidson & Humphreys 1997 and references therein). Nevertheless, having such a massive star is in fact not completely unrealistic and would have far reaching consequences if found to be true.

5. A Viable Solution to the Discrepancy

As the title suggests, there is a clear solution to the discrepancy. As the results show, the sonic point appears to be between the optical depths of ~ 1 and ~ 300 . The exact value cannot be obtained since it requires the integration outward from the photosphere, which owing to the

relatively inaccurate effective temperature and therefore opacity, yields a wide range of results. If the *mean* radiative force between the point $\eta = 1$ and the above found optical depth, is smaller than classically estimated, then a solution to the discrepancy can be found. Such a reduction in the mean radiative force is a natural result if the atmosphere is inhomogeneous.

Shaviv (1998) has shown that in an inhomogeneous atmosphere, the effective opacity used to calculate the average force is reduced relative to the effective opacity used for the radiation transfer in a homogeneous medium. The effective opacity used for the average force should be a *flux weighted* average of the opacity per unit volume³ $\chi_v \equiv \chi\rho$. Namely,

$$\chi_{eff} = \frac{\langle \chi\rho F \rangle}{\langle F \rangle \langle \rho \rangle}. \quad (8)$$

The effect is universal and arises in inhomogeneous systems that conduct heat or electricity. Extensive discussions exist in the literature under a different terminology (Isichenko 1992). The only requirement is therefore, that close to the Eddington limit the star develop inhomogeneities. The transformation from an homogeneous to an inhomogeneous atmosphere at luminosities close to but below Eddington luminosities, was recently found to take place generically even in Thomson scattering atmospheres (Shaviv 1999; Spiegel & Tao 1999; Shaviv 2000a).

It was found that two different types of instabilities arise naturally when the luminosity approaches the Eddington limit (Shaviv 2000a). One instability is of a phase transition into a *stationary* nonlinear pattern of “fingers” that facilitate the escape of the radiation. The second type of instability allows the growth of a propagating wave, from which one expects a *propagating* nonlinear pattern to form. The two possibilities are summarized in figure 2. Both instabilities bring about a reduction of the average radiative force on the matter and a significant reduction of the mass loss rate since the sonic surface can sit near (or not much below) the photosphere. In both cases, the nonlinear pattern is necessarily expected to form in the region between the radius r_{conv} at which $\eta = 1$, or in other words, that $\mathcal{L}_{conv} + \mathcal{L}_{adv}$ is large enough to have $\mathcal{L}_r \lesssim \mathcal{L}_{Edd,mod}$, and the photosphere. When the pattern is stationary, the rarefied regions have a larger than Eddington flux and the sonic surface in these regions is near r_{conv} . On the other hand, if the pattern is propagating, the flux may be larger locally than the Eddington limit but the time average of the force on a mass element is less than Eddington. Since the instability does not occur above the photosphere, it should be homogeneous and hence super-Eddington with a super-sonic flow.

Further analysis of the instabilities is needed to know which instability will dominate though it is more likely to be the phase transition since it is dynamically more important.

³When the flux is frequency dependant, a similar average should be taken in order to find the radiative force. However, one then takes a flux weighted mean over *frequency space*.

6. Summary

To summarize, the super-Eddington luminosity emitted by η -Car should have generated a much thicker wind with a sonic point placed significantly deeper than what can be directly inferred from the observations. A solution which lives in harmony with observations and theoretical modeling is a porous atmosphere, which allows more radiation to escape while exerting a smaller average force. It also means that the Eddington limit is not as destructive as one would a priori think it must be, even in a globally spherically symmetric case. Namely, all astrophysical analyses that employ the Eddington limit as a strict limit should be reconsidered carefully, even if they involve only unmagnetized Thomson scattering material. If η -Carinae could have been super-Eddington for such a long duration without “evaporating”, other systems can display a similar behavior.

REFERENCES

- Currie, D. G., Dowling, D. M., Shaya, E. J., Hester, J., Scowen, P., Groth, E. J., Lynds, R. O., & Earl, J. Jr Wide Field/Planetary Camera Instrument Definition Team, 1996, ApJ, 112, 1115
- Davidson, K. 1987, ApJ, 317, 760
- Davidson, K. & Humphreys, R. M., 1997, ARA&A, 35, 1
- Hillier, D. J. & Allen, D. A. 1992, A&A, 262, 153
- Humphreys, R. M. & Davidson, K. 1994, PASP, 106, 1025
- Iglesias, C. A. & Rogers, F. J. 1996, ApJ, 464, 943
- Isichenko, M. B. 1992, Rev. Mod. Phys., 64, 961.
- Joss, P. C., Salpeter, E. E. & Ostriker, J. P. 1973, 181, 429
- Kato, M., & Hachisu, I. 1994, ApJ, 437, 802
- Kudritzki, R. P., Pauldrach, A., Puls, J. & Abbott, D. C. 1989 A&A, 219, 205
- Mihalas D., & Weibel Mihalas B. 1984, Foundations of radiation hydrodynamics, Oxford univ. press, Oxford
- Quinn, T. & Paczynski, B. 1985, ApJ, 289, 634
- Shaviv, N. J. 1998, ApJ, 494, L193
- Shaviv, N. J. 1999, in *Variable and Non-spherical Stellar Winds in Luminous Hot Stars, IAU Colloquium 169*, ed. B. Wolf, O. Stahl, A. W. Fullerton, Springer, p. 155

Shaviv, N. J. 2000a, submitted to ApJ

Shaviv, N. J. 2000b, submitted to ApJ

Spiegel, E. A., & Tao L. 1999, Physics Reports, 311, 163

Żytkow, A. 1972, Acta Astronomica, 22, 103




Fig. 1.— The fraction $\eta = (\mathcal{L}_{adv} + \mathcal{L}_{conv})/\mathcal{L}_{crit}$ as a function of the photospheric velocity v_{ph} . A consistent wind solution requires (a) that the velocity at the photosphere v_{ph} satisfy: $v_s \leq v_{ph} \leq v_\infty$, and (b) , $\eta(v = v_s) = 1$. The thick line corresponds to the nominal observed and inferred values ($M_\star = 100 M_\odot$, $T_{eff} = 9000^\circ\text{K}$, $\dot{M} = 0.1 M_\odot/\text{yr}$, $\int \mathcal{L}_{obs} dt = 3 \times 10^{49} \text{ erg}$) while the additional lines depict the result when the values are changed to their reasonable limits (and even beyond). Clearly, no reasonable choice of parameters can result with a sonic point that is consistent with a wind solution (namely, we always find $\eta(v = v_s) \ll 1$). Basically, the discrepancy arises because the mass loss rate observed is too small to have the sonic point deep enough in the atmosphere where convection can be an efficient mean of energy transport.




Fig. 2.— The proposed atmospheric structure of η Carinae during its great eruption. A homogeneous atmosphere is unstable as a result of two generic instabilities that take place even in Thomson atmospheres when close to the Eddington limit (Shaviv 2000a). The effective opacity is therefore reduced (Shaviv 1998) and with it the average radiative force. The two panels describe the two types of possibilities for having a ‘porous’ atmosphere according to the characteristics of the instability that arises. An instability could produce a stationary pattern (first panel) if it originates from the phase transition instability and a moving pattern if it originates from the finite speed of light instability (second panel). See details in the text.